



TRIPLE CROWN®

LOW STARCH 13% Protein • 6% Fat • 18% Fiber • 13.5% NSC • 1,428 KCAL/lb.

Formulated specifically for mature horses, Triple Crown Low Starch is a pelleted low starch and sugar (NSC) feed for horses requiring a lower diet for metabolic reasons or for horse owners seeking calmer behavior out of their horses. Featuring fewer carbohydrates than hay and pasture, Triple Crown Low Starch can be used as a complete feed to replace all or part of the forage portion of the diet. Contains no corn, molasses, or alfalfa meal – ingredients believed to cause erratic behavior in horses

IDEAL FOR: Hyperactive horses, horses prone to ulcers, horses with HYPP

Guaranteed Analysis:

Crude Protein (min.)	13.00%	Iron (min.)	170 ppm
Lysine (min.)	0.70%	Manganese (min.)	100 ppm
Methionine (min.)	0.20%	Cobalt (min.)	0.29 ppm
Threonine (min.)	0.35%	Vitamin A (min.)	6,000 IU/lb.
Crude Fat (min.)	6.00%	Vitamin D (min.)	1,200 IU/lb.
Omega-3 Fatty Acids (min.)	0.30%	Vitamin E (min.)	200 IU/lb.
Omega-6 Fatty Acids (min.)	1.60%	Riboflavin (min.)	3.80 mg/lb.
Crude Fiber (max.)	18.00%	Thiamin (min.)	12.0 mg/lb.
Calcium (min.)	0.75%	Biotin (min.)	0.30 mg/lb.
Calcium (max.)	1.25%	Ascorbic Acid (min.)	45 mg/lb.
Phosphorus (min.)	0.60%	Saccharomyces Cerevisiae Yeast Culture (min.)	1.816 billion CFU/lb.
Magnesium (min.)	0.50%	Lactobacillus Acidophilus Fermentation Product (min.)	908 million CFU/lb.
Potassium (min.)	0.75%	Enterococcus Faecium Fermentation Product (min.)	908 million CFU/lb.
Copper (min.)	50 ppm	Cellulase (Trichoderma Longibrachiatum) (min.) *	0.68 million CU/lb.
Selenium (min.)	0.60 ppm	Protease (Bacillus Subtilis) (min.) **	1,430 HUT/lb.
Zinc (min.)	170 ppm		

*One cellulase unit (CU) is defined as the amount of activity that will produce a relative fluidity change of 1 (determined with a calibrated viscometer) in 5 minutes in a defined carboxymethylcellulose substrate at pH 4.5 and 40°C
 ** One hemoglobin unit (HUT) of proteolytic (protease) activity is defined as that amount of enzyme that produces, in one minute under the specified conditions, a hydrolysate whose absorbance at 275nm is the same as that of a solution containing 1.10g per mL of tyrosine in 0.006N hydrochloric acid.

Ingredients:

(Free from Restricted Ruminant Protein Products per Title 21, CFR 589.2000/2001)

Wheat Middlings, Soybean Hulls, Shredded Beet Pulp, Distillers Dried Grains, Soybean Oil, Ground Limestone, Salt, Sodium Bicarbonate, Sodium Sesquicarbonate, Stabilized Rice Bran, Calcium Carbonate, Vitamin E Supplement, Hydrated Sodium Calcium Aluminosilicate, Yeast Extract, Yeast Culture, Calcium Silicate, Sodium Aluminosilicate, Dried Enterococcus Faecium Fermentation Product, Dried Lactobacillus Acidophilus Fermentation Product, Dried Trichoderma Longibrachiatum Fermentation Extract, Dried Bacillus Subtilis Fermentation Extract, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Kelp Meal, Iron Amino Acid Complex, Magnesium Oxide, Fenugreek Seed, Lecithin, Anise Seed, Magnesium Proteinate, Selenium Yeast, Flaxseed, Monocalcium Phosphate, Dicalcium Phosphate, Lignin Sulfonate, L-Lysine, Ascorbic Acid (Source of Vitamin C), Niacin Supplement, Biotin, Vitamin A Supplement, Thiamine Mononitrate, Beta Carotene, Calcium Pantothenate, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B12 Supplement, Vitamin D3 Supplement, Choline Chloride, Menadione Sodium Bisulfite Complex, (Source of Vitamin K Activity), Folic Acid, Brewers Dried Yeast, DL-methionine, DL-methionine Hydroxy Analog, Ferrous Sulfate, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Cobalt Sulfate, Ethylenediamine Dihydroiodide.

Feeding Directions:

Triple Crown Low Starch can be fed to all horses one year of age or older. Feed enough to maintain desired body condition and energy level along with hay, and/or pasture, water and salt. If feeding less than 6 lb. per day to horses weighing 900 lb. or more, also feed additional minerals and vitamins supplied by either Triple Crown 30% or Triple Crown Lite. Triple Crown Low Starch is a complete diet and can replace all or a part of your fiber portion of your diet; however, we strongly recommend that you continue to feed a portion of your diet in long stem hay or pasture.

NOTE: Provide plenty of fresh, clean water at all times. Keep product fresh in cool, dry storage. Examine product daily. DO NOT use if old, moldy or contaminated with insects.